

Carnegie Imaging For Women

The Leading Edge in Women's Imaging

Suggested Guidelines for Fetal Echocardiography Referral

Effective January 2016

**for further information go to:

<http://www.carnegieimaging.com/wp-content/uploads/2015/10/Echo-final.pdf>

Fetal Echo Referred to Carnegie—20-22 Weeks

IVF or ICSI Pregnancies

Maternal Use of SSRI's

Pregestational Diabetes

Monchorionic Twins

Presence of Single Umbilical Artery

1st Degree Relative w/Congenital Heart Disease

Increased Nuchal Translucency (≥ 3.0 mm)

Maternal Presence SS-A or SS-B Antibodies

Maternal Lupus

Familial Inherited Disorders (ie, Marfans, 22q11.2 Deletion Syndrome)

Polyhydramnios

Maternal Seizure Disorder

Fetal Echo Referred to Pediatric Cardiology

Suspected Cardiac Abnormality During Anatomic Ultrasound

Suspected Major Fetal Anomaly

Suspected Twin-Twin Transfusion Syndrome

Known or Suspected Chromosomal Abnormalities

Teratogen Exposure during Vulnerable Period, excluding Antidepressant and Anti-epileptic Meds

Non Immune Hydrops

Prolonged Exposure to Prostaglandin Synthetase Inhibitors in 3rd Trimester (Indocin)